

SCREENING COMPONENTS	INFANCY								EARLY CHILDHOOD							LATE CHILDHOOD							ADOLESCENCE					
	NB	2 WKS	1 MOS	2 MOS	4 MOS	6 MOS	9 MOS	12 MOS	15 MOS	18 MOS	24 MOS	30 MOS	3 YRS	4 YRS	5 YRS	6 YRS	7 YRS	8 YRS	9 YRS	10 YRS	11 YRS	12 YRS	13 YRS	14 YRS	15 YRS	16 YRS	17-20 YRS (10)	
HISTORY/HISTORY UPDATE	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
VITAL SIGNS (1)																												
BLOOD PRESSURE													X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
HEIGHT/WEIGHT	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
HEAD CIRCUMFERENCE	X	X	X	X	X	X	X	X	X	X	X																	
CHEST CIRCUMFERENCE (1)																												
EYE																												
EXTERNAL EYE EXAM	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
RED REFLEX	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
LIGHT REFLEX	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
EYE MOVEMENT	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
VISUAL ACUTY (2)													X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X(11)
HEARING																												
SUBJECTIVE EXAM	X	X	X	X	X	X	X	X	X	X	X	X	X															
AUDIOMETRY														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X(11)
DENTAL																												
VISUAL DENTAL EXAM	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
DENTAL HEALTH EDUCATION	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
REFER TO DENTIST (3)											X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
UNCLOTHED PHYSICAL (4)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
DEVELOPMENTAL/ BEHAVIORAL	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
LABORATORY:																												
HEMOGLOBIN/ HEMATOCRIT (5)	**** ONCE IN THIS AGE GROUP ****								**** ONCE IN THIS AGE GROUP ****							**** ONCE IN THIS AGE GROUP ****							ONCE IN THIS AGE GROUP 1					
SICKLE CELL (6)	X																											
BLOOD LEAD(7*)								X		X	X*	X*	X*	X*	X*													
URINALYSIS (1)																												
IMMUNIZATIONS (8)																												
LEAD RISK ASSESSMENT (7)						RA	RA	RA	RA	RA	RA	RA	RA	RA	RA													
TB RISK ASSESSMENT (9)								X			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
HEALTH EDUCATION																												
ANTICIPATORY GUIDANCE	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

- At clinician's discretion.
- Should be started at age 3 years if child is cooperative. Age 3 through 7 should be done with occlusive patch. Age 8 years and older should be done with occluder.
- A child is to be referred for initial preventive dental treatment at age 2 years. If the dentist is not scheduling the child every 6 months for prophylactic and restorative dental care, the provider should refer at each visit after 3 years of age. Earlier initial dental examinations may be appropriate for some children. Subsequent examination as prescribed by dentist.
- A check for orthopedic problems is a part of the physical exam for all ages. Particular attention should be given to infants for hip displacement, club foot, and to children 8 years and older for development of Scoliosis.
- Hemoglobin/hematocrit are to be completed at least once during each age grouping: 0-14 months; 15 months - 7 years; 8 years - 13 years; 14 years - 21 years. Additional testing should be done as clinically indicated.
- Testing to be done at first visit if it has not already been done for all children of Mediterranean, African, Asian, Caribbean, South or Central American heritage.
- RA= risk assessment; BT= blood test. \* = A screening blood lead test is also required for any Medicaid eligible child 30 - 72 months of age who has not previously been screened for lead poisoning.
- Follow immunization schedule approved by ACIP, AAP, and AAFP.
- Beginning at age 12 months, TB risk assessment is to be completed at least annually. Tuberculin skin testing is based on TB risk assessment guidelines. When skin testing is needed, Pediatric Program recognizes interdermal testing but not tine testing.
- One annual visit at 17-18-19-and 20 years
- Required at age 18 only.